





IDEAL FOR:

Mature horses, performance horses, underweight horses



FEATURES:

Whole grain free, high fat & fiber, reduced starch & sugar



HELPS SUPPORT:

Poor teeth, ulcers, healthy inflammatory response, fat absorption, weight gain

NOW WITH

 EnduraCare® for healthy inflammatory response

Equ[®]Mix

- More probiotics via CLOSTAT® to support a healthy microbiome
- LYSOFORTE® for improved fat absorption and feed efficiency
- Zinpro® Performance Minerals® with higher bioavailability
- No added iron

Developed specifically for older horses, Triple Crown Senior features a beet pulp-based formula with higher nutrient and fat content to provide the digestive assistance and calories older horses require to maintain good body condition. This feed can be mixed with water to form a mash for horses that have difficulty chewing and swallowing and has enough bulk fiber to be used as a sole ration for horses unable to eat hay or pasture. It is ideal for horses of any age needing additional fiber and reduced starch and sugar levels.

14% PROTEIN • 10% FAT • 17% FIBER • 1,546 KCAL/LB

GUARANTEED ANALYSIS	
Crude Protein (min.)	14.00%
Lysine (min.)	0.80%
Methionine (min.)	0.22%
Threonine (min.)	0.66%
Tryptophan (min.)	0.18%
Leucine (min.)	0.96%
Crude Fat (min.)	10.00%
Crude Fiber (max.)	17.00%
ADF (max.)	21.00%
NDF (max.)	36.00%
Calcium (min.)	0.90%
Calcium (max.)	1.40%
Phosphorus (min.)	0.45%
Magnesium (min.)	0.37%
Potassium (min.)	1.20%
Sodium (min.)	0.40%
Sodium (max.)	0.75%
Salt (min.)	0.25%
Salt (max.)	0.50%
Omega-3 Fatty Acids (min.)	0.60%
Omega-6 Fatty Acids (min.)	3.50%
Selenium (min.)	0.30 ppm
Zinc (min.)	225 ppm
Manganese (min.)	100 ppm
Copper (min.)	50 ppm
Iron (min.)	150 ppm
Vitamin A (min.)	6,000 IU/lb.
Vitamin D min.)	1,000 IU/lb.
Vitamin E (min.)	175 IU/lb.
Riboflavin-B2 (min.)	2.50 mg/lb.
Thiamine-B1 (min.)	7.00 mg/lb.
Ascorbic Acid (min.)	45 mg/lb.
Biotin (min.)	0.20 mg/lb.
Total Direct Fed Microbials (min.)	2.50 billion CFU/lb.
Lactobacillus acidophilus Fermentation Product (min.)	590 million CFU/lb.
Enterococcus faecium Fermentation Product (min.)	590 million CFU/lb.
Bacillus subtilis Dried Fermentation Product (min.)	151 million CFU/lb.
Saccharomyces cerevisiae (min.)	1.17 billion CFU/lb.
Cellulase¹ (Trichoderma longibrachiatum) (min.)	2,137 CU/lb.
Protease ² (Bacillus licheniformis) (min.)	0.44 HUT/lb.
,, ,, ,, ,,	

1. One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulos substrate at pH 4.5 and 40°C.

INGREDIENTS

Dehydrated Alfalfa Meal, Dried Plain Beet Pulp Shreds, Soybean Hulls, Cane Molasses, Wheat Middlings, Soybean Meal, Soybean Oil, Maize Distillers Dried Grains with Solubles, Flaxseed, Heat Stabilized Rice Bran, Calcium Carbonate, Sodium Bicarbonate, Monocalcium Phosphate, Lignin Sulfonate, Magnesium Oxide, L-Lysine, Salt, Hydrated Sodium Calcium Aluminosilicate, Manganese Amino Acid Complex, Zinc Amino Acid Complex, N-Butyric Acid, Zinc Oxide, Peppermint Oil, Vitamin E Supplement, Yeast Culture, Processed Grain By-Products, Selenium Yeast, Lecithin, Copper Amino Acid Complex, DL-Methionine, Spray Dried Egg Product, Dried Kelp, Yeast Extract, Active Dry Yeast, Ascorbic Acid, Vitamin B12 Supplement, Copper Sulfate, Fenugreek, Natural and Artificial Flavors, Niacin Supplement, Magnesium Proteinate, Biotin, Maltodextrins, Dried Bacillus subtilis Fermentation Product, Dried Enterococcus faecium Fermentation Product, Dried Lactobacillus acidophilus Fermentation Product, Thiamine Mononitrate, L-Leucine, Vitamin A Supplement, β-Carotene, Pyridoxine Hydrochloride, Riboflavin Supplement, D-Calcium Pantothenate, Vitamin D3 Supplement, Choline Chloride, Zinc Sulfate, Folic Acid, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Dried Trichoderma longibrachiatum Fermentation Product, Ethylenediamine Dihydroiodide, Dried Bacillus licheniformis Fermentation Extract. Contains a source of live (viable), naturally occurring microorganisms

Contains a source of live (viable), naturally occurring microorganisms.

FEEDING DIRECTIONS

Feeding recommendations are based on average daily requirements. Horse's needs may vary based on age, work level, metabolism, weather and other factors.

Adjust the amount of Triple Crown Senior up or down as needed to maintain desired body condition. If feeding less than the amount recommended per day, supplement with Triple Crown Balancer, Balancer Gold or Diamond Balancer.

Do not feed more than 0.50% of the horse's body weight of Triple Crown Senior in a single meal, and if feeding that amount, allow 3 to 4 hours between meals.

For horses eating hay or other forage.			
Horse Weight (lb.)	Senior (lb./day)	Minimum Forage (lb./day)	
600	4-5	9-12	
900	5-10	13.5-18	
1,100	6-12	16.5-22	
1,300	7-14	19.5-26	

For horses NOT eating hay or other forage

Horse Weight (lb.)	Senior (lb./day)
600	7.5-9.5
900	10.5-14.5
1,100	13.5-17.5
1,300	16.5-20.5

Adding water to Triple Crown Senior will make an EZ Mash for horses with poor teeth, respiratory problems, or for horses prone to choking. Add water 10 to 15 minutes prior to feeding at an amount to meet the desires of the individual horse.

NOTE: Provide salt and plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects.

CAUTION: Follow label directions. Product contains Copper; do not feed to sheep.

^{2.} One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme That produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid. Estimated values determined by Equi-Analytical, Ithaca, NY.