





IDEAL FOR:

Overweight horses, ponies, miniature horses, metabolic horses



FEATURES:

Whole grain, soy & molasses free, gastric buffering, whey protein/amino acid balance, added chromium



NOW WITH

- No added iron
- Zinpro® Performance Minerals® with higher bioavailability



HELPS SUPPORT:

Balancing hay & pasture, feeding grains below recommended feeding rate, strong top line, insulin sensitivity, managing inflammation and heat stress

Triple Crown Balancer Gold is a highly fortified vitamin/mineral supplement with no soy and no molasses for horses of any age. Ideal for easy keepers, Balancer Gold can be used to balance pasture and/or hay, improve nutrition in diets when feeding below the recommended feeding rate or when mixing with whole grains (corn, oats, barley).

13.5% PROTEIN • 4.5% FAT • 15% FIBER • 1,300 KCAL/LB

GUARANTEED ANALYSIS	
Crude Protein (min.)	13.50%
Lysine (min.)	0.90%
Methionine (min.)	0.29%
Threonine (min.)	0.60%
Tryptophan (min.)	0.25%
Leucine (min.)	1.00%
Crude Fat (min.)	4.50%
Crude Fiber (max.)	15.00%
ADF (max.)	20.00%
NDF (max.)	32.00%
Calcium (min./max.)	3.00/3.50%
Phosphorus (min.)	1.50%
Magnesium (min.)	2.50%
Potassium (min.)	1.30%
Sodium (min./max.)	0.65/1.00%
Chloride (min.)	0.65%
Salt (min./max.)	0.25/1.00%
Omega-3 Fatty Acids (min.)	1.20%
Omega-6 Fatty Acids (min.)	1.00%
Selenium (added) (min.)	3.30 ppm
Selenium (max.) (total)	3.96 ppm
Zinc (min.)	1,000 ppm
Manganese (min.)	575 ppm
Copper (min.)	275 ppm
Iron (min.)	200 ppm
Chromium (max.)	6.00 ppm
Vitamin A (min.)	36,000 IU/lb.
Vitamin D min.)	6,500 IU/lb.
Vitamin E (min.)	1,000 IU/lb.
Riboflavin-B2 (min.)	18.00 mg/lb.
Thiamine-B1 (min.)	42.00 mg/lb.
Ascorbic Acid (min.)	275 mg/lb.
Biotin (min.)	1.40 mg/lb.
Total Direct Fed Microbials (min.)	18.80 billion CFU/lb.
Lactobacillus acidophilus Fermentation Product (min.)	4.54 billion CFU/lb.
Enterococcus faecium Fermentation Product (min.)	4.54 billion CFU/lb.
Bacillus subtilis Dried Fermentation Product (min.)	664 million CFU/lb.
Saccharomyces cerevisiae (min.)	9.06 billion CFU/lb.
Cellulase¹ (Trichoderma longibrachiatum) (min.)	16,436 CU/lb.
Protease ² (Bacillus licheniformis) (min.)	3.40 HUT/lb.
(

^{1.} One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C.

INGREDIENTS

(Free from Restricted Ruminant Protein Products per Title 21, CFR 589.2000/2001) Alfalfa Meal, Wheat Middlings, Ground Flaxseed Meal, Monocalcium Phosphate, Magnesium Oxide, Dried Whey, Calcium Carbonate, Lignosulfonate, Calcite, Manganese Amino Acid Complex, Zinc Amino Acid Complex, Selenium Yeast, Butvric Acid. Zinc Oxide, Peppermint Essential Oil, Potassium Chloride, Saccharomyces cerevisiae Yeast Grown On A Media Of Sucrose and Cane Molasses, Processed Grain By-Products, Salt, L-Lysine, Vitamin E Supplement, Copper Amino Acid Complex, Hydrated Sodium Calcium Aluminosilicate, Chromium Propionate, Leucine, Dried Kelp, L-Threonine, Yeast Extract, Active Dry Yeast, DL-Methionine, L-Tryptophan, Ascorbic Acid (Source of Vitamin C), Vitamin B12 Supplement, Ground Fenugreek, Natural & Artificial Flavors, Magnesium Proteinate, Niacin Supplement, Biotin, Copper Sulfate, Maltodextrins, Dried Bacillus subtilis Fermentation Product, Dried Enterococcus faecium Fermentation Product, Dried Lactobacillus acidophilus Fermentation Product, Thiamine Mononitrate, Vitamin A Supplement, Beta-Carotene, Pyridoxine Hydrochloride, Riboflavin Supplement, D-Calcium Pantothenate, Vitamin D3 Supplement, Choline Chloride, Dried Trichoderma longibrachiatum Fermentation Product, Folic Acid, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Zinc Sulfate, Ethylenediamine Dihydroiodide, Dried Bacillus licheniformis Fermentation Extract.

Contains a source of live (viable), naturally occurring microorganisms.

FEEDING DIRECTIONS

For easy keeping, mature horses.			
Horse weight (lb.)	Balancing pasture or hay-only diets	Balancing unfortified grain diets*	Balancing diets when feeding less than the recommended amount of fortified feed
600 - 900	0.5 - 0.75 lb./day	0.5 - 0.75 lb./day	0.25 - 0.50 lb./day
900 - 1,100	0.75 - 1 lb./day	0.75 - 1 lb./day	0.50 - 0.75 lb./day
1,100 - 1,300	1 - 1.5 lb./day	1 - 1.5 lb./day	0.75 - 1 lb./day
1,300	1.5 lb./day	1.5 lb./day	1 lb./day
Weanlings 400 - 600	0.5 - 0.75 lb./day	0.5 - 0.75 lb./day	0.5 - 0.75 lb./day
Yearlings 700 - 800	1 lb./day	1 lb./day	1 lb./day

*provides appropriate vitamin/mineral fortification according to body weight when feeding otherwise unfortified sources of calories such as oats, beet pulp or rice bran. For desired body condition, adjust the calorie source (oats, beet pulp, rice bran) up or down to adjust caloric intake.

OTHER CONSIDERATIONS

Feeding recommendations assume a horse is consuming a minimum of 1.50 to 2% of its body weight in hay or pasture and are based on average daily requirements. Horse's needs may vary based on age, work level, metabolism, weather and other factors.

NOTE: Provide plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects

CAUTION: Follow feeding guidelines. Chromium supplementation should not exceed 4 mg per day. Feeding added Selenium at levels in excess of 0.30 ppm in the total diet is prohibited. Product contains added Copper; do not feed to sheep.

^{2.} One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid. Estimated values determined by Equi-Analytical, Ithaca, NY.