




BALANCER

PELLET

50 LBS
(22.7 KG)



IDEAL FOR:

Overweight horses, broodmares, growing horses, metabolic horses

FEATURES:

Whole grain free, low starch & sugar, low feeding rate, added chromium

HELPS SUPPORT:

Balancing hay & pasture, feeding grains below recommended feeding rate, insulin sensitivity, managing inflammation and heat stress

IMMUNITY
Equi-Mix
GUT HEALTH

NOW WITH

- More probiotics via CLOSTAT® to support a healthy microbiome
- Zinpro® Performance Minerals® with higher bioavailability
- No added iron

While originally formulated for growing horses and broodmares, Triple Crown Balancer is a highly fortified, low sugar and starch pelleted supplement, ideal for easy keepers and horses with metabolic disease and insulin resistance. This can be used to balance pasture and/or hay and improve nutrition in diets when feeding below the recommended feeding rate and/or mixing with whole grains.

30% PROTEIN • 3% FAT • 5% FIBER • 1,266 KCAL/LB

GUARANTEED ANALYSIS

Crude Protein (min.)	30.00%
Lysine (min.)	2.00%
Methionine (min.)	0.62%
Threonine (min.)	1.35%
Tryptophan (min.)	0.44%
Leucine (min.)	2.25%
Crude Fat (min.)	3.00%
Crude Fiber (max.)	5.00%
ADF (max.)	7.00%
NDF (max.)	13.00%
Calcium (min./max.)	4.00/5.00%
Phosphorus (min.)	1.50%
Magnesium (min.)	2.40%
Potassium (min.)	1.30%
Sodium (min./max.)	0.50/1.00%
Salt (min./max.)	0.95/1.45%
Omega-3 Fatty Acids (min.)	0.40%
Omega-6 Fatty Acids (min.)	1.10%
Selenium (min./max.)	3.00/4.00 ppm
Zinc (min.)	1,000 ppm
Manganese (min.)	575 ppm
Copper (min.)	275 ppm
Iron (min.)	150 ppm
Chromium (max.)	6.00 ppm
Vitamin A (min.)	36,000 IU/lb.
Vitamin D min.)	6,500 IU/lb.
Vitamin E (min.)	1,000 IU/lb.
Riboflavin-B2 (min.)	16.00 mg/lb.
Thiamine-B1 (min.)	42.00 mg/lb.
Ascorbic Acid (min.)	275 mg/lb.
Biotin (min.)	1.40 mg/lb.
Total Direct Fed Microbials (min.)	18.68 billion CFU/lb.
Lactobacillus acidophilus Fermentation Product (min.)	4.54 billion CFU/lb.
Enterococcus faecium Fermentation Product (min.)	4.54 billion CFU/lb.
Bacillus subtilis Dried Fermentation Product (min.)	598 million CFU/lb.
Saccharomyces cerevisiae (min.)	9.00 billion CFU/lb.
Cellulase ¹ (Trichoderma longibrachiatum) (min.)	16,436 CU/lb.
Protease ² (Bacillus licheniformis) (min.)	3.40 HUT/lb.

1. One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C.

2. One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.09g per mL of tyrosine in 0.006N hydrochloric acid. Estimated values determined by Equi-Analytical, Ithaca, NY.

INGREDIENTS

Soybean Meal, Wheat Middlings, Calcium Carbonate, Monocalcium Phosphate, Magnesium Oxide, Lignin Sulfonate, Flaxseed, Salt, Soybean Oil, Sodium Bicarbonate, Cane Molasses, Lecithin, Manganese Amino Acid Complex, Rice Bran, Zinc Amino Acid Complex, Selenium Yeast, N-Butyric Acid, Zinc Oxide, Peppermint Oil, Yeast Culture, Processed Grain By-Products, Vitamin E Supplement, L-Lysine, DL-Methionine, Copper Amino Acid Complex, Hydrated Sodium Calcium Aluminosilicate, Chromium Propionate, Dried Kelp, Yeast Extract, Active Dry Yeast, Ascorbic Acid, Vitamin B12 Supplement, Fenugreek, Natural and Artificial Flavors, Magnesium Proteinate, Niacin Supplement, Biotin, Copper Sulfate, Dried Enterococcus faecium Fermentation Product, Dried Lactobacillus acidophilus Fermentation Product, Dried Bacillus subtilis Fermentation Product, Leucine, Thiamine Mononitrate, Vitamin A Supplement, β-Carotene, Pyridoxine Hydrochloride, Riboflavin Supplement, D-Calcium Pantothenate, Vitamin D3 Supplement, Choline Chloride, Dried Trichoderma longibrachiatum Fermentation Product, Folic Acid, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Zinc Sulfate, Ethylenediamine Dihydroiodide, Dried Bacillus licheniformis Fermentation Extract.

Contains a source of live (viable), naturally occurring microorganisms.

FEEDING DIRECTIONS

Horse Weight (lb.)	Amount of Balancer (lb./day)		
	Balancing Pasture or Hay-Only Diets	Balancing Unfortified Grain Diets	Balancing Diets when Feeding Less than the Recommended Amount of Fortified Feed
600	0.5 - 0.75	0.5 - 0.75	0.25 - 0.50
900	0.75 - 1	0.75 - 1	0.50 - 0.75
1,100	1 - 1.5	1 - 1.5	0.75 - 1
1,300	1.5	1.5	1
Weanlings 400 - 600	0.5 - 0.75	0.5 - 0.75	0.5 - 0.75
Yearlings 700 - 800	1	1	1

The above feeding recommendations assume a horse is consuming a minimum of 1.50 to 2% of its body weight in hay or pasture for appropriate vitamin/mineral fortification according to body weight. Horse's needs may vary based on age, work level, metabolism, weather and other factors.

For desired body condition, adjust the amount of forage or unfortified concentrate inclusion (oats, corn, barley, rice bran, beet pulp) up or down to adjust caloric intake. If more than the recommended amount of Triple Crown Balancer is needed to maintain body condition, consider switching to Triple Crown Lite or adding a Triple Crown performance or senior feed.

NOTE: Provide salt and plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects.

CAUTION: Follow label directions. Chromium supplementation should not exceed 4 mg per day. Feeding added Selenium at levels in excess of 0.30 ppm in the total diet is prohibited. Triple Crown Balancer does not exceed an intake of 3 mg of added Selenium per head per day when fed as directed. Product contains added Copper; do not feed to sheep.

Triple Crown® and EquiMix® are registered trademarks of Triple Crown Nutrition, Inc. 315 Lake Street East, Suite #300, Wayzata, MN 55391; CLOSTAT® is a registered trademark of Kemin Industries, Inc., U.S.A.; Zinpro® and Zinpro Performance Minerals® are registered trademarks of Zinpro Corporation.