



IDEAL FOR:

Mature horses, metabolic horses, underweight horses



FEATURES:

Whole grain & molasses free, low starch & sugar, balanced vitamin & mineral fortification



+

HELPS SUPPORT:

Poor teeth, hay shortages, poor hay quality, improved insulin senstivity

Triple Crown Safe Starch® Fortified Forage is a complete forage-based diet for mature horses, providing an alternative to grain-based diets

PROTEIN 11% • FAT 6% • FIBER 28% • 1,100 KCAL/LB

GUARANTEED ANALYSIS	
Crude Protein (min.)	11.00%
Lysine (min.)	0.35%
Methionine (min.)	0.10%
Threonine (min.)	0.30%
Tryptophan (min.)	0.07%
Leucine (min.)	0.38%
Crude Fat (min.)	6.00%
Crude Fiber (max.)	28.00%
ADF (max.)	30.00%
NDF (max.)	46.00%
Calcium (min.)	0.40%
Calcium (max.)	0.90%
Phosphorus (min.)	0.20%
Magnesium (min.)	0.25%
Potassium (min.)	1.80%
Selenium (added) (min.)	0.30 ppm
Zinc (min.)	100 ppm
Manganese (min.)	100 ppm
Copper (min.)	35 ppm
Iron (min.)	150 ppm
Chromium (min.)	0.44 ppm
Vitamin A (min.)	4,000 IU/lb.
Vitamin D (min.)	700 IU/lb.
Vitamin E (min.)	100 IU/lb.
Ascorbic Acid (min.)	30 mg/lb.
Biotin (min.)	0.15 mg/lb.
Total Direct Fed Microbials (min.)	1.75 billion CFU/lb.
Lactobacillus acidophilus Fermentation Product (min.)	454 million CFU/lb.
Enterococcus faecium Fermentation Product (min.)	454 million CFU/lb.
Bacillus subtilis Dried Fermentation Product (min.)	66 million CFU/lb.
Saccharomyces cerevisiae (min.)	908 million CFU/lb.
Cellulase¹ (Trichoderma longibrachiatum Fermentation Extract) (min.)	0.34 million CU/lb.
Protease ^{2, 3} (Bacillus licheniformis) (min.)	715 HUT/lb. or PC/lb.

¹One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C"

²One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid.

 3 One bacterial protease unit (PC) is defined as the quantity of enzyme that produces the equivalent of 1.5µg/mL of L-tyrosine per minute under assay conditions of pH 7.0 and 37°C.

INGREDIENTS

(Free from Restricted Ruminant Protein Products per Title 21, CFR 589.2000/2001) Orchard Hay, Timothy Hay, Glycerin, Wheat Middlings, Soybean Oil, Dehulled Soybean Meal, Calcium Carbonate, Ground Soybean Hulls (0.70%), Magnesium Oxide, Monocalcium Phosphate, Manganese Amino Acid Complex, Butyric Acid, Zinc Oxide, Peppermint Essential Oil, Zinc Amino Acid Complex, Saccharomyces cerevisiae Yeast Grown On A Media Of Sucrose, Cane Molasses, Processed Grain By-Products, Vitamin E Supplement, Sodium Bicarbonate, Salt, Lignosulfonate, Copper Amino Acid Complex, Hydrolyzed Yeast, Cereal Distillers Dried Solubles, Fat Product Feed Grade, Monosodium Phosphate, Dried Kelp, Yeast Extract, L-Lysine, Active Dry Yeast, Ascorbic Acid, DL-Methionine, Vitamin B12 Supplement, Ground Fenugreek, Natural & Artificial Flavors, Hydrated Sodium Calcium Aluminosilicate, Selenium Yeast, Chromium Proprionate, Magnesium Proteinate, Niacin Supplement, Biotin, Copper Sulfate, Maltodextrins, Dried Bacillus subtilis Fermentation Product, Dried Enterococcus faecium Fermentation Product, Dried Lactobacillus acidophilus Fermentation Product. Leucine, Thiamine Mononitrate, Vitamin A Supplement, Beta-Carotene, Pyridoxine Hydrochloride, Riboflavin Supplement, D-Calcium Pantothenate, Vitamin D3 Supplement, Choline Chloride, Dried Trichoderma longibrachiatum Fermentation Product Folic Acid Menadione Sodium Risulfite Complex Zinc Sulfate, Ethylenediamine Dihydroiodide, Dried Bacillus licheniformis Fermentation Extract.

Contains a source of live (viable), naturally occurring microorganisms.

FEEDING DIRECTIONS	
TO SUPPLEMENT HAY/PASTURE	
1,000 lb. horse	4 - 6 lb. per day
TO REPLACE FEED/GRAIN	
1,000 lb. horse	10 lb. per day
TO USE AS A COMPLETE DIET	
1,000 lb. horse	15 - 20 lb. per day

OTHER CONSIDERATIONS

Product can be mixed with water to aid in hydration. Product may be mixed with the grain portion of the diet to slow rate of intake.

When feeding as a replacement to feed/grain, should be fed in combination with a minimum of 0.50% of body weight in hay/pasture.

Provide salt and plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects.

CAUTION STATEMENT

Follow feeding guidelines. Chromium supplementation should not exceed 4 mg per day. Feeding added selenium at levels in excess of 0.30 ppm in the total diet is prohibited. Product contains added copper; do not feed to sheep.