



SAFE STARCH® FORAGE

FORTIFIED FORAGE

40 LBS
(18.4 KG)



IDEAL FOR:

Mature horses, metabolic horses, underweight horses



FEATURES:

Whole grain & molasses free, low starch & sugar, balanced vitamin & mineral fortification



HELPS SUPPORT:

Poor teeth, hay shortages, poor hay quality, insulin sensitivity



NOW WITH
KemTRACE®
Chromium

Triple Crown Safe Starch® Fortified Forage is a complete forage-based diet for mature horses, providing an alternative to grain-based diets

11% PROTEIN • 6% FAT • 28% FIBER • 1,100 KCAL/LB

GUARANTEED ANALYSIS

Crude Protein (min.)	11.00%
Lysine (min.)	0.35%
Methionine (min.)	0.10%
Threonine (min.)	0.30%
Tryptophan (min.)	0.07%
Leucine (min.)	0.38%
Crude Fat (min.)	6.00%
Crude Fiber (max.)	28.00%
ADF (max.)	30.00%
NDF (max.)	46.00%
Calcium (min.)	0.40%
Calcium (max.)	0.90%
Phosphorus (min.)	0.20%
Magnesium (min.)	0.25%
Potassium (min.)	1.80%
Selenium (added) (min.)	0.30 ppm
Zinc (min.)	100 ppm
Manganese (min.)	100 ppm
Copper (min.)	35 ppm
Iron (min.)	150 ppm
Chromium (max.)	0.44 ppm
Vitamin A (min.)	4,000 IU/lb.
Vitamin D (min.)	700 IU/lb.
Vitamin E (min.)	100 IU/lb.
Ascorbic Acid (min.)	30 mg/lb.
Biotin (min.)	0.15 mg/lb.
Total Direct Fed Microbials (min.)	1.88 billion CFU/lb.
Lactobacillus acidophilus Fermentation Product (min.)	454 million CFU/lb.
Enterococcus faecium Fermentation Product (min.)	454 million CFU/lb.
Bacillus subtilis Dried Fermentation Product (min.)	66.40 million CFU/lb.
Saccharomyces cerevisiae (min.)	906 million CFU/lb.
Cellulase ¹ (Trichoderma longibrachiatum) (min.)	1,644 CU/lb.
Protease ² (Bacillus licheniformis) (min.)	0.34 HUT/lb.

¹One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C

²One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid.

INGREDIENTS

(Free from Restricted Ruminant Protein Products per Title 21, CFR 589.2000/2001) Orchard Hay, Timothy Hay, Glycerin, Wheat Middlings, Soybean Oil, Dehulled Soybean Meal, Calcium Carbonate, Ground Soybean Hulls (0.70%), Magnesium Oxide, Monocalcium Phosphate, Manganese Amino Acid Complex, Butyric Acid, Zinc Oxide, Peppermint Essential Oil, Zinc Amino Acid Complex, Saccharomyces cerevisiae Yeast Grown On A Media Of Sucrose and Cane Molasses, Processed Grain By-Products, Vitamin E Supplement, Sodium Bicarbonate, Salt, Lignosulfonate, Copper Amino Acid Complex, Hydrolyzed Yeast, Maize Distillers Dried Grains with Solubles, Dried Seaweed Meal, Monosodium Phosphate, Dried Kelp, Yeast Extract, L-Lysine, Active Dry Yeast, Ascorbic Acid, DL-Methionine, Vitamin B12 Supplement, Ground Fenugreek, Natural & Artificial Flavors, Hydrated Sodium Calcium Aluminosilicate, Selenium Yeast, Chromium Propionate, Magnesium Proteinates, Niacin Supplement, Biotin, Copper Sulfate, Maltodextrins, Dried Bacillus subtilis Fermentation Product, Dried Enterococcus faecium Fermentation Product, Dried Lactobacillus acidophilus Fermentation Product, Leucine, Thiamine Mononitrate, Vitamin A Supplement, Beta-Carotene, Pyridoxine Hydrochloride, Riboflavin Supplement, D-Calcium Pantothenate, Vitamin D3 Supplement, Choline Chloride, Dried Trichoderma longibrachiatum Fermentation Product, Folic Acid, Menadione Sodium Bisulfite Complex, Zinc Sulfate, Ethylenediamine Dihydroiodide, Dried Bacillus licheniformis Fermentation Extract.

Contains a source of live (viable), naturally occurring microorganisms.

FEEDING DIRECTIONS

TO SUPPLEMENT HAY/PASTURE

1,100 lb. horse 4 – 6 lb. per day

TO REPLACE FEED/GRAIN

1,100 lb. horse 10 lb. per day

TO USE AS A COMPLETE DIET

1,100 lb. horse 15 – 20 lb. per day

OTHER CONSIDERATIONS

Product can be mixed with water to aid in hydration. Product may be mixed with the grain portion of the diet to slow rate of intake.

When feeding as a replacement to feed/grain, should be fed in combination with a minimum of 0.50% of body weight in hay/pasture.

Provide salt and plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects.

CAUTION STATEMENT

Follow feeding guidelines. Chromium supplementation should not exceed 4 mg per day. Feeding added Selenium at levels in excess of 0.30 ppm in the total diet is prohibited. Product contains added Copper; do not feed to sheep.