SENIOR ACTIVE+

EXTRUDED PELLET

40 LBS
(18.14 KG)



IDEAL FOR:
Active, older horses

FEATURES:
Whole grain free, high fat & fiber, extruded for enhanced digestibilty, gastric buffering

HELPS SUPPORT:
Ulcers, poor teeth

TSC TRACTOR SUPPLY CO

EquiMix IMMUNITY GUT HEALTH

NOW WITH
- No added iron
- Zinpro® Performance Minerals® with higher bioavailability

Developed for active, older horses, Triple Crown Senior Active+ is extruded for enhanced digestibility while providing gastric buffering which is ideal for performance horses or those under stress. This feed can be can be mixed with water to form a mash for horses that have difficulty chewing or swallowing.

PROTEIN 14% • FAT 12% • FIBER 16% • 1,535 KCAL/LB

GUARANTEED ANALYSIS

Crude Protein (min.)	14.00%
Lysine (min.)	0.95%
Methionine (min.)	0.30%
Threonine (min.)	0.80%
Tryptophan (min.)	0.15%
Leucine (min.)	0.97%
Crude Fat (min.)	12.00%
Crude Fiber (max.)	16.00%
ADF (max.)	22.00%
NDF (max.)	34.00%
Calcium (min.)	0.75%
Calcium (max.)	1.25%
Phosphorus (min.)	0.65%
Salt (min.)	0.25%
Salt (max.)	0.75%
Sodium (min.)	0.25%
Sodium (max.)	0.75%
Magnesium (min.)	0.55%
Potassium (min.)	0.99%
Chloride	0.50%
Omega-3 Fatty Acids (min.)	1.20%
Omega-6 Fatty Acids (min.)	5.50%
Copper (min.)	50 ppm
Selenium (min./max.)	0.50/0.90 ppm
Zinc (min.)	220 ppm
Manganese (min.)	150 ppm
Iron (min.)	150 ppm
Vitamin A (min.)	5,930 IU/lb.
Vitamin D3 (min.)	1,125 IU/lb.
Vitamin E (min.)	170 IU/lb.
Riboflavin-B2 (min.)	2.50 mg/lb.
Thiamin-B1 (min.)	7.00 mg/lb.
Biotin (min.)	0.20 mg/lb.
Ascorbic Acid (min.)	40 mg/lb.
Total Direct Fed Microbials (min.)	4.14 billion CFU/lb.
Saccharomyces cerevisiae (min.)	1.81 billion CFU/lb.
Lactobacillus acidophilus (min.)	1.13 billion CFU/lb.
Enterococcus faecium (min.)	1.13 billion CFU/lb.
Bacillus subtilis (min.)	71.90 million CFU/lb.
Cellulase[1] (Trichoderma longibrachiatum) (min.)	3,287 CU/lb.
Protease[2] (Bacillus licheniformis) (min.)	0.68 HUT/lb.

1. One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C.

2. One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid. Estimated values determined by Equi-Analytical, Ithaca, NY.

INGREDIENTS

Wheat Middlings, Soybean Hulls, Dehydrated Alfalfa Meal, Soybean Oil, Maize Distillers Dried Grains with Solubles, Dried Plain Beet Pulp, Steamed Whole Oats, Flaxseed, Stabilized Rice Bran, Calcite, Soybean Meal, Monocalcium Phosphate, Sodium Bicarbonate, Salt, L-Lysine, Magnesium Oxide, L-Threonine, Hydrolyzed Yeast, Dried Seaweed Meal, Manganese Amino Acid Complex, Dried Enterococcus faecium Fermentation Product, Dried Lactobacillus acidophilus Fermentation Product, DL-Methionine, Yeast Culture, Natural and Artificial Flavors, Zinc Amino Acid Complex, Selenium Yeast, Vitamin E Supplement, Calcium Carbonate, L-Leucine, Hydrated Sodium Calcium Aluminosilicate, Copper Amino Acid Complex, Propyl Gallate, Tocopherols, Citric Acid, Dried Kelp, Yeast Extract, Zinc Sulfate, Active Dry Yeast, Ascorbic Acid, Fenugreek, Vitamin B12 Supplement, Magnesium Proteinate, Niacin Supplement, Biotin, Copper Sulfate, Dried Bacillus subtilis Fermentation Product, Thiamine Mononitrate, Vitamin A Supplement, β-Carotene, Pyridoxine Hydrochloride, Riboflavin Supplement, D-Calcium Pantothenate, Vitamin D3 Supplement, Choline Chloride, Dried Trichoderma longibrachiatum Fermentation Product, Folic Acid, Menadione Sodium Bisulfite Complex (Source of Vitamin K Activity), Ethylenediamine Dihydroiodide, Dried Bacillus licheniformis Fermentation Extract

Contains a source of live (viable), naturally occurring microorganisms.

FEEDING DIRECTIONS

FEED FOR OPTIMAL CONDITION:
Feeding recommendations are based on average daily requirements. Horse's needs may vary based on age, work level, metabolism, weather and other factors. Adjust the amount of Triple Crown Senior Active+ up or down as needed to maintain desired body condition. Triple Crown Senior Active+ must be fed with adequate forage.

If feeding less than the recommended amount per day, ensure your horse isn't missing essential vitamins and minerals by supplementing with Triple Crown Balancer, Triple Crown Balancer Gold or Triple Crown Diamond Balancer.

Do not feed more than 0.50% of the horse's body weight of Triple Crown Senior Active+ in a single meal, and if feeding that amount, allow 3 to 4 hours between meals.

Horse Weight (lb.)	Senior Active+ (lb./day)	Minimum Forage (lb./day)
600	3-4	9-12
900	4-9	13.5-18
1,100	5-11	16.5-22
1,300	6-13	19.5-26

NOTE:
Provide salt and plenty of fresh, clean water at all times.
Keep product fresh in cool, dry storage. Examine product daily.
DO NOT use if old, moldy or contaminated with insects.

CAUTION:
Follow label directions.Feeding added Selenium at levels in excess of 0.30 ppm in the total diet is prohibited.
Triple Crown Senior Active+ does not exceed an intake of 3 mg of added Selenium per head per day when fed as directed.
Product contains Copper; do not feed to sheep.