



# SENIOR ACTIVE+

EXTRUDED PELLET

40 LBS  
(18.14 KG)



**IDEAL FOR:**  
Active, older horses



**FEATURES:**  
Whole grain free, high fat & fiber, extruded for enhanced digestibility, gastric buffering



**HELPS SUPPORT:**  
Ulcers, poor teeth

Developed for active, older horses, Triple Crown Senior Active+ is extruded for enhanced digestibility while providing gastric buffering which is ideal for performance horses or those under stress. This feed can be mixed with water to form a mash for horses that have difficulty chewing or swallowing.

**PROTEIN 14% • FAT 12% • FIBER 16% • 1,535 KCAL/LB**



## GUARANTEED ANALYSIS

Crude Protein (min.)	14.00%
Lysine (min.)	0.95%
Methionine (min.)	0.30%
Threonine (min.)	0.80%
Tryptophan (min.)	0.15%
Leucine (min.)	0.97%
Crude Fat (min.)	12.00%
Crude Fiber (max.)	16.00%
ADF (max.)	22.00%
NDF (max.)	34.00%
Calcium (min.)	0.75%
Calcium (max.)	1.25%
Phosphorus (min.)	0.65%
Magnesium (min.)	0.55%
Potassium (min.)	0.99%
Sodium (min.)	0.25%
Sodium (max.)	0.75%
Chloride	0.50%
Salt (min.)	0.25%
Salt (max.)	0.75%
Omega-3 Fatty Acids (min.)	1.20%
Omega-6 Fatty Acids (min.)	5.50%
Selenium (min./max.)	0.60/1.10 ppm
Zinc (min.)	220 ppm
Manganese (min.)	95 ppm
Cobalt (min.)	0.20 ppm
Copper (min.)	50 ppm
Iron (min.)	200 ppm
Vitamin A (min.)	5,930 IU/lb.
Vitamin D3 (min.)	1,125 IU/lb.
Vitamin E (min.)	170 IU/lb.
Riboflavin-B2 (min.)	2.50 mg/lb.
Thiamin-B1 (min.)	7.00 mg/lb.
Ascorbic Acid (min.)	40 mg/lb.
Biotin (min.)	0.20 mg/lb.
Total Direct Fed Microbials (min.)	4.05 billion CFU/lb.
Lactobacillus Acidophilus Fermentation Product (min.)	1.13 billion CFU/lb.
Enterococcus Faecium Fermentation Product (min.)	1.13 billion CFU/lb.
Bacillus Subtilis Dried Fermentation Product (min.)	27 million CFU/lb.
Saccharomyces Cerevisiae Yeast Culture (min.)	1.77 billion CFU/lb.
Cellulase <sup>1</sup> (Trichoderma Longibrachiatum) (min.)	0.68 million CU/lb.
Protease <sup>2</sup> (Bacillus Subtilis) (min.)	1,430 HUT/lb.

1. One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C.

2. One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid. Estimated values determined by Equi-Analytical, Ithaca, NY.

## INGREDIENTS

(Free from Restricted Ruminant Protein Products per Title 21, CFR 589.2000/2001)

Wheat Middlings, Soybean Hulls, Dehydrated Alfalfa Meal, Soybean Oil, Dried Plain Beet Pulp, Distiller's Dried Grains with Solubles, Oat Groats, Whole Flaxseed, Stabilized Rice Bran, Soybean Meal, Sodium Bicarbonate, Calcite, Calcium Carbonate, Monocalcium Phosphate, Salt, Magnesium Oxide, L-Lysine Hydrochloride, L-Tryptophan, DL-Methionine, L-Threonine, L-Leucine, Dextrose, Dried Enterococcus Faecium Fermentation Product, Dried Lactobacillus Acidophilus Fermentation Product, Bacillus Subtilis, Hydrolyzed Yeast, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Hydrated Sodium Calcium Aluminosilicate, Yeast Extract, Kelp Meal, Dried Yeast, Lecithin, Fenugreek Seed, Magnesium Proteinate, Selenium Yeast, Vitamin E Supplement, Ascorbic Acid, Niacin Supplement, Biotin, Thiamine Mononitrate, Vitamin A Supplement, Vitamin B12 Supplement, Beta Carotene, Pyridoxine Hydrochloride, Riboflavin Supplement, Calcium Pantothenate, Vitamin D3 Supplement, Choline Chloride, Folic Acid, Menadione Sodium Bisulfite Complex, Ferrous Sulfate, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Cobalt Proteinate, Ethylenediamine Dihydroiodide, Cobalt Sulfate, and Natural and Artificial Flavors.

## FEEDING DIRECTIONS

Start with the amount of feed recommended for the given weight and work level of the horse; adjust amount being fed over time to achieve desired body condition. If feeding less than 5 lb. per day per 1,000 lb. of body weight, add Triple Crown Balancer to provide essential vitamins and minerals at a level that meets your horse's daily requirements. Please contact Triple Crown with any questions you might have about your horse's feeding program.

800 LB. HORSE		
Work level	Feeding rate (lb.)	Feeding notes
Light	3.5 - 4.5 per day	To ensure optimal fiber consumption, feed a minimum of 12 lb. of hay per day
Moderate	4.5 - 5.5 per day	
Heavy	5.5 - 6.5 per day	
1,000 LB. HORSE		
Work level	Feeding rate (lb.)	Feeding notes
Light	5.0 - 6.0 per day	To ensure optimal fiber consumption, feed a minimum of 15 lb. of hay per day
Moderate	5.5 - 6.5 per day	
Heavy	7.0 - 8.0 per day	
1,200 LB. HORSE		
Work level	Feeding rate (lb.)	Feeding notes
Light	6.5 - 7.5 per day	To ensure optimal fiber consumption, feed a minimum of 18 lb. of hay per day
Moderate	7.0 - 8.0 per day	
Heavy	8.0 - 9.0 per day	
1,400 LB. HORSE		
Work level	Feeding rate (lb.)	Feeding notes
Light	8.0 - 9.0 per day	To ensure optimal fiber consumption, feed a minimum of 21 lb. of hay per day
Moderate	8.5 - 9.5 per day	
Heavy	9.5 - 10.5 per day	

**NOTE:** Provide plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects.