




BALANCER

PELLET

50 LBS
(22.7 KG)



IDEAL FOR:

Overweight horses, broodmares, growing horses, metabolic horses



FEATURES:

Whole grain free, low starch & sugar, low feeding rate



HELPS SUPPORT:

Balancing hay & pasture, feeding grains below recommended feeding rate, managing inflammation and heat stress

NOW WITH
CHROMIUM!

While originally formulated for growing horses and broodmares, Triple Crown Balancer is a highly fortified, low sugar and starch pelleted supplement, ideal for easy keepers and horses with metabolic disease and insulin resistance. This can be used to balance pasture and/or hay and improve nutrition in diets when feeding below the recommended feeding rate and/or mixing with whole grains.

PROTEIN 30% • FAT 3% • FIBER 5% • 1,266 KCAL/LB

GUARANTEED ANALYSIS

Crude Protein (min.)	30.00%
Lysine (min.)	2.00%
Methionine (min.)	0.62%
Threonine (min.)	1.35%
Tryptophan (min.)	0.44%
Leucine (min.)	2.25%
Crude Fat (min.)	3.00%
Crude Fiber (max.)	5.00%
ADF (max.)	7.00%
NDF (max.)	13.00%
Calcium (min./max.)	4.00/5.00%
Phosphorus (min.)	1.50%
Magnesium (min.)	2.40%
Potassium (min.)	1.30%
Sodium (min./max.)	0.50/1.00%
Salt (min./max.)	0.95/1.45%
Omega-3 Fatty Acids (min.)	0.40%
Omega-6 Fatty Acids (min.)	1.10%
Selenium (min./max.)	3.40/4.40 ppm
Zinc (min.)	1,000 ppm
Manganese (min.)	575 ppm
Cobalt (min.)	2.10 ppm
Copper (min.)	300 ppm
Iron (min.)	500 ppm
Chromium (max.)	8.80 ppm
Vitamin A (min.)	36,000 IU/lb.
Vitamin D min.)	6,000 IU/lb.
Vitamin E (min.)	1,000 IU/lb.
Riboflavin-B2 (min.)	16.00 mg/lb.
Thiamine-B1 (min.)	42.00 mg/lb.
Ascorbic Acid (min.)	275 mg/lb.
Biotin (min.)	1.40 mg/lb.
Total Direct Fed Microbials (min.)	17.97 billion CFU/lb.
Lactobacillus Acidophilus Fermentation Product (min.)	4.54 billion CFU/lb.
Enterococcus Faecium Fermentation Product (min.)	4.54 billion CFU/lb.
Bacillus Subtilis Dried Fermentation Product (min.)	169 million CFU/lb.
Saccharomyces Cerevisiae (min.)	8.69 billion CFU/lb.
Cellulase ¹ (Trichoderma Longibrachiatum) (min.)	850 CU/lb.
Protease ² (Bacillus Licheniformis) (min.)	3.08 HUT/lb.

INGREDIENTS

(Free from Restricted Ruminant Protein Products per Title 21, CFR 589.2000/2001) Soybean Meal, Wheat Middlings, Calcium Carbonate, Monocalcium Phosphate, Magnesium Oxide, Heat Stabilized Rice Bran, Soybean Oil, Molasses, Ground Flaxseed, Salt, Lignosulfonate, Sodium Bicarbonate, Selenium Yeast, Butyric Acid, Zinc Oxide, Peppermint Essential Oil, Saccharomyces Cerevisiae Yeast Extract, Saccharomyces Cerevisiae Active Dry Yeast, Hydrated Sodium Calcium Aluminosilicate, Dried Enterococcus Faecium Fermentation Product, Dried Lactobacillus Acidophilus Fermentation Product, Dried Trichoderma Longibrachiatum Fermentation Product, Dried Bacillus Licheniformis Fermentation Extract, Saccharomyces Cerevisiae Yeast Culture, Vitamin E Supplement, DL-Methionine, Chromium Propionate, L-Threonine, L-Lysine, Dried Kelp, Zinc Sulfate, Manganese Sulfate, Zinc Hydroxychloride, Ascorbic Acid, L-Leucine, Manganese Hydroxychloride, Vitamin B12 Supplement, Lecithin, Natural & Artificial Flavors, Fenugreek Seed, Magnesium Proteinates, Niacin Supplement, Copper Hydroxychloride, Copper Sulfate, Biotin, Thiamine Mononitrate, Vitamin A Supplement, Beta-Carotene, Dried Bacillus Subtilis Fermentation Product, Cobalt Proteinates, Pyridoxine Hydrochloride, Riboflavin Supplement, D-Calcium Pantothenate, Cobalt Sulfate, Vitamin D3 Supplement, Choline Chloride, Folic Acid, Menadione Sodium Bisulfite Complex, Ethylenediamine Dihydroiodide.

FEEDING DIRECTIONS

TO BALANCE PASTURE AND/OR HAY³

Ponies: 500 - 700 lb.	0.5 - 0.75 lb. per day
Weanlings: 400 - 600 lb.	0.5 - 0.75 lb. per day
Yearlings: 700 - 800 lb.	1 lb. per day
Horses: 800 - 1,100 lb.	1 - 1.50 lb. per day
Broodmares and Breeding Stallions	1.5 - 2 lb. per day
Horses in moderate-to-high levels of activity	1.5 - 2 lb. per day

TO BALANCE DIETS WHEN FEEDING BELOW RECOMMENDED FEEDING RATE

Ponies: 500 - 700 lb.	0.25 - 0.5 lb. per day
Weanlings: 400 - 600 lb.	0.5 - 0.75 lb. per day
Yearlings: 700 - 800 lb.	1 lb. per day
Horses: 800 - 1,100 lb.	0.5 - 1 lb. per day
Broodmares and Breeding Stallions	1 lb. per day
Horses in moderate-to-high levels of activity	1 lb. per day

TO BALANCE GRAINS⁴

Ponies: 500 - 700 lb.	0.5 - 0.75 lb. per day
Horses: 800 - 1,100 lb.	1 - 1.5 lb. per day
Broodmares and Breeding Stallions	1.5 - 2 lb. per day
Horses in moderate-to-high levels of activity	1.5 - 2 lb. per day

OTHER CONSIDERATIONS

The above feeding recommendations assume a horse is consuming a minimum of 1.5 to 2% of its body weight in hay or pasture.

Provide salt and plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects.

1. One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C.
 2. One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid. Estimated values determined by Equi-Analytical, Ithaca, NY.
 3. Assuming a horse is consuming a minimum of 1.5 to 2% of its body weight in hay or pasture.
 4. The feeding recommendations provide for appropriate vitamin/mineral fortification according to body weight. For desired body condition, adjust grain inclusion (oats, corn, barley) to provide additional calories.