



IDEAL FOR:

Mature horses, performance horses, underweight horses







FEATURES:

Whole grain free, gastric buffering, antioxidant support



HELPS SUPPORT:

Ulcers, stamina, immune function, oxidative stress

Triple Crown Diamond Perform is a high fat, high fiber pellet with antioxidants, designed to meet the caloric needs of performance horses and others requiring additional calories while promoting gut and immune health.

12.5% PROTEIN • 12% FAT • 15% FIBER

GUARANTEED ANALYSIS	
Crude Protein (min.)	12.50%
Lysine (min.)	0.80%
Methionine (min.)	0.28%
Threonine (min.)	0.55%
Tryptophan (min.)	0.20%
Leucine (min.)	0.90%
Crude Fat (min.)	12.00%
Crude Fiber (max.)	15.00%
ADF (max.)	19.00%
NDF (max.)	34.00%
Calcium (min./max.)	1.00/1.50%
Phosphorus (min.)	0.60%
Magnesium (min.)	0.45%
Potassium (min.)	1.00%
Sodium (min./max.)	0.50/1.00%
Chloride (min.)	0.45%
Salt (min./max.)	0.60/1.00%
Omega-3 Fatty Acids (min.)	2.00%
Omega-6 Fatty Acids (min.)	3.90%
Selenium (min./max.)	0.50/1.00 ppm
Zinc (min.)	220 ppm
Manganese (min.)	95 ppm
Copper (min.)	50 ppm
Iron (min.)	60 ppm
Vitamin A (min.)	5,900 IU/lb.
Vitamin D min.)	1,100 IU/lb.
Vitamin E (min.)	210 IU/lb.
Riboflavin-B2 (min.)	2.50 mg/lb.
Thiamine-B1 (min.)	7.00 mg/lb.
Ascorbic Acid (min.)	250 mg/lb.
Biotin (min.)	0.22 mg/lb.
Total Direct Fed Microbials (min.)	3.62 billion CFU/lb.
Lactobacillus acidophilus Fermentation Product (min.)	907 million CFU/lb.
Enterococcus faecium Fermentation Product (min.)	907 million CFU/lb.
Bacillus subtilis Dried Fermentation Product (min.)	66.25 million CFU/lb.
Saccharomyces Cerevisiae (min.)	1.74 billion CFU/lb.
Cellulase¹ (Trichoderma longibrachiatum) (min.)	170 CU/lb.
Protease ^{2,3} (Bacillus licheniformis) (min.)	0.62 HUT/lb. or PC/lb.

1. One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C

INGREDIENTS

(Free from Restricted Ruminant Protein Products per Title 21, CFR 589.2000/2001) Wheat Middlings, Dehydrated Alfalfa Meal, Ground Flaxseed Soybean Oil, Ground Beet Pulp, Molasses, Heat Stabilized Rice Bran, Lignosulfonate, Maize Distillers Dried Grains, Ground Oats, Calcite, Salt, Monocalcium Phosphate, Sodium Bicarbonate, Magnesium Oxide, Calcium Carbonate, L-Lysine, L-Tryptophan, Butyric Acid, Zinc Oxide, Peppermint Essential Oil, L-Threonine, DL-Methionine Ascorbic Acid, Selenium Yeast, Vitamin E Supplement, Manganese Amino Acid Complex, Saccharomyces cerevisiae Yeast Culture, Zinc Amino Acid Complex, Hydrated Sodium Calcium Aluminosilicate, Dried Kelp, Copper Amino Acid Complex, Saccharomyces cerevisiae Yeast Extract, Saccharomyces cerevisiae Active Dry Yeast, Vitamin B12 Supplement, Ground Fenugreek, Natural & Artificial Flavors, Magnesium Proteinate, Copper Amino Acid Complex, Niacin Supplement, Biotin, Dried Enterococcus faecium Fermentation Product, Dried Lactobacillus acidophilus Fermentation Product, Copper Sulfate, Leucine, Dried Bacillus subtilis Fermentation Product, Thiamine Mononitrate, Vitamin A Supplement, Beta-Carotene, Pyridoxine Hydrochloride, Riboflavin Supplement, D-Calcium Pantothenate, Vitamin D3 Supplement, Choline Chloride, Dried Trichoderma longibrachiatum Fermentation Product, Folic Acid, Menadione Sodium Bisulfite Complex, Zinc Sulfate, Ethylenediamine Dihydrojodide, Dried Bacillus licheniformis Fermentation Extract.

Contains a source of live (viable), naturally occurring microorganisms.

FEEDING DIRECTIONS

Feeding recommendations are based on average daily requirements. Horses' needs may vary based on age, work level, metabolism, weather and other factors. Adjust the amount of Triple Crown Diamond Perform, up or down, as needed to maintain desired body condition. If less than the amount recommended is fed per day, supplement with Triple Crown Diamond Balancer.

Do not feed more than 0.50% of the horse's body weight of Triple Crown Diamond Perform in a single meal. Allow 3 to 4 hours between meals

NOTE: Provide salt and plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage Examine product daily. DO NOT use if old, moldy or contaminated with insects.

CAUTION STATEMENT: Follow label directions. Feeding added Selenium at levels in excess of 0.30 ppm in the total diet is prohibited.

600 LB. HURSE			
Work Level	Feeding Rate/Day	Feeding Notes	
Idle	2 lb.	Feed a minimum of 9-12 lb. of forage per day	
In Work	3-4 lb.		
900 LB. HORSE			
Work Level	Feeding Rate/Day	Feeding Notes	
Idle	3 lb.	Feed a minimum of 13.5-18 lb. of forage	
In Work	4-9 lb.	per day	
1,100 LB. HORSE			
Work Level	Feeding Rate/Day	Feeding Notes	
Idle	3.5 lb.	Feed a minimum of	
In Work	5-11 lb.	16.5-22 lb. of forage per day	
1,300 LB. HORSE			
Work Level	Feeding Rate/Day	Feeding Notes	
Idle	4 lb.	Feed a minimum of 19.5-26 lb. of forage per day	
In Work	6-13 lb.		

^{2.} One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.10g per mL of tyrosine in 0.006N hydrochloric acid.

Estimated values determined by Equi-Analytical, Ithaca, NY.

3. One bacterial protease unit (PC) is defined as the quantity of enzyme that produces the equivalent of 1.5µg/mL of L-tyrosine per minute under assay conditions of pH 7.0 and 37°C.