




# DIAMOND SENIOR

TEXTURED

50 LBS  
(22.7 KG)



**IDEAL FOR:**  
Mature horses, performance horses



**FEATURES:**  
Whole grain free, gastric buffering, antioxidant support, EZ Mash



**HELPS SUPPORT:**  
Ulcers, poor teeth, immune function, oxidative stress

found exclusively at  
**TSC TRACTOR SUPPLY CO**



Triple Crown Diamond Senior is a beet pulp-based, high fat, high fiber feed that provides antioxidant support for immune function. It is perfect for senior horses and others that need a higher nutrient and calorie intake to maintain good body condition.

**14.5% PROTEIN • 12.5% FAT • 18% FIBER • 1,800 KCAL/LB.**

## GUARANTEED ANALYSIS

Crude Protein (min.)	14.50%
Lysine (min.)	0.90%
Methionine (min.)	0.25%
Threonine (min.)	0.70%
Tryptophan (min.)	0.20%
Leucine (min.)	1.00%
Crude Fat (min.)	12.50%
Crude Fiber (max.)	18.00%
ADF (max.)	22.00%
NDF (max.)	37.00%
Calcium (min./max.)	0.90/1.40%
Phosphorus (min.)	0.55%
Magnesium (min.)	0.40%
Potassium (min.)	1.00%
Sodium (min./max.)	0.50/1.00%
Chloride (min.)	0.50%
Salt (min./max.)	0.40/0.90%
Omega-3 Fatty Acids (min.)	1.10%
Omega-6 Fatty Acids (min.)	5.00%
Selenium (min.)	0.30 ppm
Zinc (min.)	260 ppm
Manganese (min.)	140 ppm
Copper (min.)	70 ppm
Iron (min.)	175 ppm
Vitamin A (min.)	6,000 IU/lb.
Vitamin D min.)	1,000 IU/lb.
Vitamin E (min.)	200 IU/lb.
Riboflavin-B2 (min.)	3.20 mg/lb.
Thiamine-B1 (min.)	8.00 mg/lb.
Ascorbic Acid (min.)	250 mg/lb.
Biotin (min.)	0.25 mg/lb.
Total Direct Fed Microbials (min.)	3.87 billion CFU/lb.
Lactobacillus acidophilus Fermentation Product (min.)	930 million CFU/lb.
Enterococcus faecium Fermentation Product (min.)	930 million CFU/lb.
Bacillus subtilis Dried Fermentation Product (min.)	158 million CFU/lb.
Saccharomyces cerevisiae (min.)	1.85 billion CFU/lb.
Cellulase <sup>1</sup> (Trichoderma longibrachiatum) (min.)	3,353 CU/lb.
Protease <sup>2,3</sup> (Bacillus licheniformis) (min.)	0.69 HUT/lb.

## INGREDIENTS

Dried Beet Pulp Shreds, Dehydrated Alfalfa Meal, Wheat Middlings, Soybean Hulls, Soybean Oil, Dehulled Soybean Meal, Maize Distillers Dried Grains, Molasses, Flaxseed, Stabilized Rice Bran, Sodium Bicarbonate, Monocalcium Phosphate, Calcite, Calcium Carbonate, Salt, Lignin Sulfonate, Magnesium Oxide, L-Lysine, L-Threonine, N-Butyric Acid, Zinc Oxide, Peppermint Oil, DL-Methionine, Selenium Yeast, Ascorbic Acid, L-Tryptophan, Propionic Acid (a Preservative), L-Leucine, Manganese Amino Acid Complex, Zinc Amino Acid Complex, Saccharomyces cerevisiae Yeast Culture, Vitamin E Supplement, Copper Amino Acid Complex, Hydrated Sodium Calcium Aluminosilicate, Dried Kelp, Saccharomyces cerevisiae Yeast Extract, Saccharomyces cerevisiae Active Dry Yeast, Vitamin B12 Supplement, Manganese sulfate, Fenugreek, Natural and Artificial Flavors, Magnesium Proteinate, Niacin Supplement, Copper Sulfate, Biotin, Dried Bacillus subtilis Fermentation Product, Dried Enterococcus faecium Fermentation Product, Thiamine Mononitrate, Vitamin A Supplement, β-Carotene, Pyridoxine Hydrochloride, Riboflavin Supplement, D-Calcium Pantothenate, Vitamin D3 Supplement, Choline Chloride, Dried Trichoderma longibrachiatum Fermentation Product, Zinc Sulfate, Folic Acid, Menadione Sodium Bisulfite Complex, Ethylenediamine Dihydroiodide, Dried Bacillus licheniformis Fermentation Extract.

Contains a source of live (viable), naturally occurring microorganisms.

## FEEDING DIRECTIONS

Feeding recommendations assume a horse is consuming a minimum of 1.50 to 2% of its body weight in hay or pasture and are based on average daily requirements. Horses' needs may vary based on age, work level, metabolism, weather and other factors. Adjust the amount of Triple Crown Diamond Senior, up or down, as needed to maintain desired body condition. If less than the amount recommended is fed per day, supplement with Triple Crown Diamond Balancer or Balancer.

Do not feed more than 0.50% of the horse's body weight of Triple Crown Diamond Senior in a single meal. Allow 3 to 4 hours between meals.

**NOTE:** Provide salt and plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects.

### CAUTION STATEMENT:

Follow label directions. Product contains added Copper; do not feed to sheep.

### 600 LB. HORSE

Feeding Rate/Day	Feeding Notes
3-4 lb.	Feed a minimum of 9-12 lb. of forage per day

### 900 LB. HORSE

4-9 lb.	Feed a minimum of 13.5-18 lb. of forage per day
---------	---

### 1,100 LB. HORSE

5-11 lb.	Feed a minimum of 16.5-22 lb. of forage per day
----------	---

### 1,300 LB. HORSE

6-13 lb.	Feed a minimum of 19.5-26 lb. of forage per day
----------	---

1. One cellulase unit (CU) is defined as the amount of activity that will produce a relative fluidity change of 1 (determined with a calibrated viscometer) in 5 minutes in a defined carboxymethylcellulose substrate at pH 4.5 and 40°C.

2. One hemoglobin unit (HUT) of proteolytic (protease) activity is defined as that amount of enzyme that produces, in one minute under the specified conditions, a hydrolysate whose absorbance at 275nm is the same as that of a solution containing 1.0g per mL of tyrosine in 0.006N hydrochloric acid. Estimated values determined by Equi-Analytical, Ithaca, NY.